

Elanco US Inc.

Product Bulletin

Permethrin™ Fly & Louse Dust

EPA Reg. No. 11556-168

For control of adult (fly) *Cochliomyia hominivorax* (New World screwworm)

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATES OF AL, AR, CA, FL, GA, LA, MS, NC, SC, and TN

FIFRA Section 2(ee) Recommendation

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or accepted by the U.S. Environmental Protection Agency.

It is a violation of federal law to use this product in a manner inconsistent with its label.

This product bulletin must be in the possession of the user at the time of pesticide application.

All applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label must be followed.

This product has not been tested specifically for efficacy against New World screwworm (NWS, adults or larvae); however, applied as part of an integrated pest management plan, it may have preventative action against adult NWS (flies).

No single therapy should be assumed to be 100% effective. Implementation of a multi-faceted program is considered critical.

APPLICATION DIRECTIONS

Refer to the full EPA-registered label for Permethrin™ Fly & Louse Dust for complete use directions and restrictions including dosage, concentration, and application frequency requirements for use on cattle and horses.

TARGET ANIMAL	APPLICATION INSTRUCTIONS
Beef and Dairy Cattle	Dust Bag Application: Place contents of this package in any commercially available dust bag, suspend bags in areas frequently by cattle or in gateways or lanes through which the animals pass daily for water, feed or minerals. Bags may also be suspended in loafing sheds or in front of protected mineral feeders. For lactating dairy cows, bags may be suspended in the exit through which cows leave the milking barn. In all cases, the bags should be adjusted so that the bottom of the bag will hang four to six inches below the topline of the cattle. For reduction of face flies, the bags must be located so the animals will be forced to use them daily and hung at the height to ensure that the faces of the cattle will be dusted.
	Direct Application: Apply 2.0 ounces of dust per animal by shaker can over the head, neck shoulders, back and tailhead. Repeat as necessary.
Swine	Direct Application: Apply not more than 1 ounce* (3 level tablespoons) per head as a uniform coat to the head, shoulders and back by use of a shaker can. Repeat as necessary, but not more often than once every ten days. In severe infestations, both individual animals and the bedding may be treated as directed above.
Do not ship animals for slaughter within 5 days of last treatment.	
*Note: 1 ounce of this material equals approximately three level tablespoonfuls.	

IMPORTANT: Read the entire Directions for Use and the Limited Warranty and Limitation of Damages before using this product.

Registrant: Elanco US Inc., 450 Elanco Circle, Indianapolis, Indiana 46221

Version 06/12/2026
Expiration 05/01/2029