

Elanco US Inc.

Product Bulletin

Permethrin™ CDS Pour-on Insecticide

EPA Reg. No. 11556-170

For control of adult (fly) *Cochliomyia hominivorax* (New World screwworm)

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATES OF AL, AR, CA, FL, GA, LA, MS, NC, SC, AND TN

FIFRA Section 2(ee) Recommendation

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or accepted by the U.S. Environmental Protection Agency.

It is a violation of federal law to use this product in a manner inconsistent with its label.

This product bulletin must be in the possession of the user at the time of pesticide application.

All applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label must be followed.

This product has not been tested specifically for efficacy against New World screwworm (adults or larvae); however, applied as part of an integrated pest management plan, it may have preventative action against adult NWS (flies).

No single therapy should be assumed to be 100% effective. Implementation of a multi-faceted program is considered critical.

APPLICATION DIRECTIONS

Refer to the full EPA-registered label for Permethrin™ CDS Pour-on Insecticide for complete use directions and restrictions including dosage, concentration, and application frequency requirements.

TARGET	APPLICATION INSTRUCTIONS
Beef and Dairy Cattle (including Lactating) and Calves	For lice and severe horn fly control apply 2.0 mL per 100 lbs. body wt. of animal, up to a maximum of 20 mL per animal. <i>Pour-on:</i> Pour correct dose along back and down face. <i>Ready-to-use Spray:</i> Use undiluted in a mist sprayer to apply correct dose. Apply directly to neck, face, back, legs and ears. <i>Backrubber:</i> Mix 64 mL (2.1 oz.) per gallon of mineral oil. Keep rubbing device charged. Results improved by daily forced use.
Sheep and Lambs	<i>Pour-on:</i> Pour along the back. Apply 1 mL per 50 lb. body weight of animal, up to maximum of 12 mL for any one animal. For optimum control, all animals in the flock should be treated after shearing. <i>Ready-to-use Spray:</i> Use undiluted in a mist sprayer to apply 4 mL of product per animal to the ventral (underside) of sheep.
Horses and Foals	<i>Pour-on:</i> Do not use this application method on foals. After riding or exercise, pour 8 mL to 16 mL per animal along back and down the face of MATURE horses, being careful to avoid the eyes. Repeat treatment as needed, but not more than once every two weeks. <i>Wipe-on:</i> Apply 8 mL to 16 mL. Dampen an applicator mitt, cloth or toweling (Turkish). Rub over hair with special attention to the legs, shoulders, neck and facial areas where flies tend to congregate. Avoid the eyes. Repeat treatment as needed, but not more than once every two weeks. <i>Ready-to-use Spray:</i> Use after riding or exercising. Apply 8 mL to 16 mL undiluted in a mist sprayer per animal. Apply directly to ticks. Repeat treatment as needed, but not more than once every two weeks.
Horse, Beef, Dairy, Swine, Sheep and Poultry Premises, Animal Hospital Pens and Kennels	<i>Ready-to-use Spot or Premise Spray:</i> Use undiluted in a mist sprayer. Apply directly to surface to leave a residual insecticidal coating, paying particular attention to areas where insects crawl or alight. 500 mL will treat approximately 7,200 square feet.

IMPORTANT: Read the entire Directions for Use and the Limited Warranty and Limitation of Damages before using this product.

Registrant: Elanco US Inc., 450 Elanco Circle, Indianapolis, Indiana 46221

Version 06/13/2026
Expiration 05/01/2029